Dental Journal

Majalah Kedokteran Gigi

Vol. 42. No. 4 October-December 2009

Research Report

The ability of IgY to recognize surface proteins of *Streptococcus* mutans

Basri A. Gani¹, Santi Chismirina¹, Zinatul Hayati², Endang Winiati B³, Boy M. Bachtiar³, and I. Wayan T. Wibawan⁴

- ¹ Department of Oral Biology, Faculty of Dentistry, Syiah Kuala University, Banda Aceh
- ² Department of Microbiology, Faculty of Dentistry, Syiah Kuala University, Banda Aceh
- ³ Department of Oral Biology, Faculty of Dentistry, Indonesia University, Jakarta
- ⁴ Department of Bacteriology and Immunology, Vertirinary Faculty, Pertanian Bogor Institute, Bogor

ABSTRACT

Background: Streptococcus mutans are gram positive bacteria classified into viridians group, and have a role in pathogenesis of dental caries. It's adhesion to the tooth surface is mediated by cell surface proteins, which interact with specific receptor located in tooth pellicle. Glucan binding protein, Glukosyltransferase, and antigen I/II are basic proteins of S. mutans, which have a role in initiating the interaction. A previous study showed that chicken's IgY can interfere the interaction. Purpose: The objective of this study was to assess the ability of IgY in recognizing the surface molecule of Streptococcus mutans expressed by various serotypes (c, d, e, f) and a strain derived from IPB, Bogor. Method: Western blot was used as a method to determine such capability. Result: The result showed that IgY has a potency to recognize antigen I/II, but not the other proteins on the cell surface of all bacteria tested. Conclusion: The ability of IgY to bind the surface protein, antigen I/II, indicates that this avian antibody could be used as a candidate for anti-adhesion in preventing dental caries.

Key words: IgY, Streptococcus mutans, adhesion, surface proteins, and dental caries

Correspondence: Basri A. Gani, c/o: Prodi Kedokteran Gigi, Fakultas Kedokteran, Universitas Syiah Kuala Darussalam Banda Aceh 23111. E-mail: basriunoe@yahoo.com

INTRODUCTION

Streptococcus mutans (S. mutans) are bacteria that have an important role in the pathogenesis of dental caries. Among bacterium species in oral cavity, S. mutans are known as one of bacteria that have recently been discussed because of its ability in forming extracellular polysaccaride and plaque. These bacteria were isolated from dental plaque for the first time by Clarke in 1924. These bacteria are classified into the Monera kingdom, Firmicutes division, Bacilli class, Lactobacillus ordo, Streptococcaceae family, Streptococcus genus, and Streptococcus mutant species. In 1890, Miller formulated chemo-parasitic theory of dental caries, known as hypothesis of non-specific plaque describing a process from decalcification process of enamel to the forming of dental caries as the cumulative effect of acid produced by bacteria of dental plaque.

Molecularly, the surface proteins of *S. mutans* that are commonly involved in the process of dental caries are

Glucan binding protein (Gbp) and Antigen I/II (Ag I/II). 4,5 Besides that, *S. mutans* also express molecules as enzyme in the process of carbohydrate fermentation that has a role in the activity of *S. mutans*, such as *Glucosyltransferase* (Gtf), Dextranase (Dex), and Fruktosiltranferase (Ftf). Each of those three enzymes then breaks sucrose in order to form glukan, dextran, and fruktan. Actually, there are other proteins that have a role in supplying energy reverse of *S. mutans* in order to be able in oral cavity, such as Dextranase A (DexA), Dextranase B (DexB), Fruktanase, and Dlt1-4 (protein intracellular cells of *S. mutans*). 6,7

Furthermore, there are two approaches for preventing caries, which are *host* aspects like diet control, clean control of oral cavity, fluorides, and agent aspects obtained from both active and passive immunizations. The use of mammalian Immunoglobulin G (IgG) in active and passive immunizations can prevent the adhesion of *S. mutans* on dental pellicle by recognizing cell surface proteins of *S. mutans*, especially Gbp, Gtf, and AgI/II, so the antibody

can be used for preventing dental caries. Besides mammalian antibody like IgG, Immunoglobulin Y (IgY) can also be used as immune-prophylaxis matter for preventing dental caries through its use as complement materials of tooth paste used for decreasing the percentage of *S. mutans* in human saliva. Tests for the IgG in passive immunization actually has been done, but the producing process of IgG has many technique limitations, is not economics, and can hurt animal that produce it during blood taking process. Based on the phenomenon, passive immunization with antibody (IgY) specifically located in egg yolk was used to substitute the role of IgG in preventing the interaction of adhesive molecule of *S. mutans* on dental surface. 10,11

Chicken's egg is the source of IgY that potentially can be used as immunotherapy materials since it is more resistant to temperature (60-65° C) and pH alteration, and does not cause cross-reaction with structural component of tissue and protein in mammalians compared with IgG. These characters indicate that IgY can be applied to diagnose and prevent disease. 12 Based on some aspects in dentistry, IgY anti S. mutans actually can be used to decrease the frequency of caries in the experiments on caries. 13 For instance, based on the result of the experiment in which rats are used as testing animal, it was known that IgY has a potency in preventing the activity of Gtf and GbpB, so it can present colonization of S. mutans in dental pellicle, but the serotype is not known. 14,15 Besides that, the reference about the ability of IgY relating with antigen I/II of S. mutans with many serotypes is still not known since protein has an important role in the adhesion of S. mutans to dental pellicle.

Therefore, this research was aimed to analyze the potency of IgY anti *S. mutans* in recognizing cell surface proteins of *S. mutans* with serotype c, d, e, and f. The significance of this experiment is to inform both the potency of IgY located in egg yolk of chicken as anti-*S. mutans*, especially for the researchers and pharmacy companies, and the possibility of IgY as passive immunization materials.

MATERIAL AND METHOD

This research was an experimental laboratory conducted in Molecular Laboratory in Faculty of Dentistry-UI and Faculty of Veterinary-IPB. Specific IgY anti *S. mutans* produced by Faculty of Veterinary, Pertanian Bogor Institute (FKH-IPB) was used as detector of bacteria from 4 serotypes, *S. mutans* serotype c (mutans Xc), *S. mutans* serotype d (sobrinus OMZ176), *S. mutans* serotype e (mutans LM7), and *S. mutans* serotype f (mutans OMZ175). Those four serotypes of *S. mutan* were laboratory strain, obtained from Professor Yamashita, Department of Preventive Dentistry, Faculty of Dentistry, Kyushu University, Japan. *S. mutans* strain derived from IPB was used as positive control.

In order to obtain antigen, each serotype of *S. mutans* was cultured in the selective solid media of Trypticase Soy

with Sucrose and Bacitracin (TYS20B) and incubated for 12–72 hours at the temperature of 37° C. Next, one colony was taken usingh *oase* after being cultivated in the water media of *Trypticase Soy Broth* (TSB), and incubated for 24–72 hours at the temperature of 37° C in microaerophilic setting. Afterwards, *whole cell S. mutans* were centrifuged at 3000 rpm for 10 minutes. Sediment obtained was added with 200 μ l lisozim, and incubated at ice temperature for about 5 minutes before centrifuged again at 3000 rpm for 5 minutes. Next, the sediment, as *S. mutans* antigen, was re-suspended in 500 μ l Phosphate Buffer Saline (PBS) and added with 12.5% Sodium Dodecyl Sulphate (SDS) for about 65 μ l.

Before the profile of cell proteins of S. mutans with serotype c, d, e, f, and a strain derived from IPB with western blot method was detected, the level of protein was determined by using Bradford (Bio-Rad) method in order to make the number of cell proteins of S. mutans that was analyzed have the same level of protein for each. S. mutans cells that have already been prepared with lisozim then were put into *Elisa plate* wells for about 160 µl $(10 \,\mu l \text{ sample} + 150 \,\mu l \text{ PBS})$. Next, other same Elisa plate wells were given with Bovine Serum Antibody (BSA) as standard proteins about $160 \mu l$ ($10 \mu l$ BSA + $150 \mu l$ PBS). Either sample or BSA was added with 40 μ l protein assay (Bradford), and then re-suspended by using multi-chanel pipet and incubated at the room temperature for 1 hour. The concentration of proteins was measured by using Elisa reader based on Optical Density with 655 nm wave length.

Western blot technique was used in order to detect the reactivity of IgY. Based on the method conducted by Yera et al., ¹⁶ the method will be as the following; SDS PAGE. The prepared collector gel was put into SDS tank that has already been filled with reservoir running buffer stock (Bio-Rad). Next, separator gel was put into $20\,\mu$ l sample and $5\,\mu$ l standard protein (Invitrogen) in different wells as the mass indicator of the sample protein molecule. Then, protein was separated through 2 phases. First, the sample and marker proteins in the wells were collected near separator gel by setting up the electricity vertically at $100\,\text{mA}$, $100\,\text{volt}$, and $16\,\text{watt}$ for $30\,\text{minutes}$. Second, the proteins in separator gel were separated by setting up the electricity at $100\,\text{mA}$, $150\,\text{voltage}$, $25\,\text{watt}$, for $80\,\text{minutes}$.

The next step was electrotransfer of antigen protein. First, gel containing antigen protein was put onto blotting nitrocellulose paper, and then was immersed in buffer transfer blot for about 2 minutes inside glass container. At the same time, transferring cassette that has already been halved was prepared. Second, each of them was put in foam pad (sponge), and then filter papers of mini trans-blot were put on them. Third, gel was put on one of them with blotting nitrocellulose on it. Afterward, the surface part was leveled up in order not to cause any bubble. Then, the transferring cassette was girded again and put into electrophoresis tank. Finally, electro-transfer was used with 100 Voltage of electricity for 1 hour.

Detection of Antigen Protein with IgYwas done. First, Blotting *nitrocellulose* (Bio-Rad) paper containing antigen was incubated in 20 ml 5% milk on sheker for about 1 hour at the temperature of 30° C. Next, it was washed 3 times with PBS (Sigma) for about 5 minutes. IgY in 10 ml 5% non-fat milk with the concentration 1:2000 was poured into the surface of blotting papers that have already been washed and incubated on shaker for about 1 hour at 30° C. Then, they were washed 4 times with PBS for 5 minutes. In order to visualize the interaction between IgY and targeted antigen protein, those blotting papers must be added with anti IgY antibody labeling peroxides (Horse rabbit peroxide anti-Chicken/Turkey IgG) with 1:2000 concentration that have already diluted with 10 ml 5% milk. Next, they were incubated on shaker for 1 hour at the temperature of 30° C. After that, those blotting papers were washed 4 times with PBS for 5 minutes. To visualize the binding between IgY and targeted antigen, finally, developing reagent was used.

RESULT

In this research, the series of working procedures for determining the profile and reactivity of IgY towards cell proteins of *S. mutans* with serotype c, d, e, and f, were conducted by using Western Blot technique. *Streptococus mutans* that have already been extrated with lisozim was analyzed by SDS PAGE method and colored with *Commasie blue* in order to analyze the profile of *S. mutans* proteins visualised in Figure 1. Paralelly, blotting of IgY anti *S. mutans* was conducted by using Western Blot technique in order to analyze cell proteins of *S. mutans* binding with IgY (Figure 2).




Figure 1. The protein profile of *S. mutans* colored with Commassie blue using SDS-PAGE method *S. mutans* in line 1, 2, and 3 showed similar protein ribbon profile, while, *S. mutans* in line 4 and 5 shows the different protein ribbon from the previous one.




Figure 2. The surface proteins of *S. mutans* recognized by IgY anti *S. mutans*, using Western Blot method (Figure 2), showed similar protein ribbon with Molecule Mass (MM) about 188 kDa for all of those five isolates of *S. mutans*. Number 1–5 show that the well with 20 µl antigen extracted with lisozim from whole cels of *S. mutans*. Line 1 (*S. mutans* Isolate FKH IPB), Line 2–4, each of which shows *S. mutans* with serotype f, serotype e, serotype d, and serotype c. M = prestaining protein marker, Kda = kilo Dalton, and the number under it showed the molecule mass of proteins.

DISCUSSION

Chicken's egg yolk is considered as the food component which also contain specific antibody towards antigen inducing the respond of the chicken's immune system. The production of IgY from chicken's egg actually can give more advantages than the production of imunoglobulin from other animals, like mammals. ¹⁴ IgY, moreover, can be used to detect some viral brands through Elisa test, immunodiffusion, and immunofluorescence since the score of its isoelectrics is lower than human's IgG. ¹⁷ IgY can also be used to detect immunoglobulin in animal serum. ¹⁴

The technology for producing IgY as an alternative antibody substituting antibody produced by mammalian animals, moreover, is considered as one of reasons for using it in immunotherapy and immunopropilaxis. ¹⁸ in dentistry, IgY can actually be used as antibody of anti *S. mutans*. ^{15,19} Thus, IgY, as anti adhesion, is expected to have potency to recognize the surface proteins causing the dental caries. In this case, the surface proteins of *S. mutans* commonly reported as the initiator of the adhesion of bacteria to dental pellicle are Gbp, Gtf, and Ag I/II. ²⁰

The choosing of whole cell *S. mutans* as the vaccine materials used for producing chicken's antibody was aimed to expose many kinds of surface epitope used for producing many kinds of paratop with many binding sites in hypervariabel regio, thus, IgY produced can interact with epitop of the surface proteins of *S. mutans* with high aviditas.²¹ It means that the binding of the surface proteins of *S. mutans* by IgY antibody can decrease the character of

S. mutans pathogens since they can restrain the adhesion of those bacteria in the *host* dental surface, so it can possibly restrain the process of caries.²²

The result from this research can give important information about the potency of IgY as anti surface proteins of *S. mutans*. This research may oven answer whether IgY produced by FKH IPB can interact with high avidities towards the surface proteins of *S. mutans*. Therefore, in order to answer the question, Western Blot technique then was used in this research.

Based on the analysis result of protein profile with commasie blue and the use of IgY anti *S. mutans* with Western Blot technique (Figure 1 and 2), the surface proteins of *S. mutans* that can be recognized by IgY were proteins with Molecule Mass (BM) about 188 kDa, while other proteins was not detected by Western Blot technique. Idone⁷ and Matos-Graner²³ report that proteins, about 180-190 kDa, are considered as Ag I/II proteins of *S. mutans*, which means that IgY used in this research can specifically recognize antigens I/II considered as the surface proteins expressed by those serotypes of *S. mutans*, meanwhile Gtf and Gbp can not be detected. The reason of this IgY potency is because fragment antigen binding (FAB) of IgY can recognize the surface proteins of *S. mutans*.²⁴

Thus, this result supports the assumption that chicken immunized with whole cell S. mutans can produce specific IgY towards Ag I/II expressed by those four serotypes of S. mutans. The potency of IgY is possibly related with the higher concentration of antibody in egg yolk.²⁵ However, this hypothesis still needs further researches. One of them conducted by Nikki, 26 shows that besides Ag I/II has 190 kDa Molecule Mass, Ag I/II also locates in the position that is relatively higher than the cell surface of S. mutans compared with Gtf and Gbp. Therefore, Ag I/II become the surface proteins that are more dominant as the initiator of adhering to dental pellicle. By using of IgY with water soluble fraction (WSF) and immune-blotting technique, Wibawan¹⁴ and Smith²⁴ reports that IgY WSF can detect 59 kDa proteins considered as the surface proteins of S. mutans (GbpA), nevertheless, IgY only shows the weak reactivation towards Pac proteins classified into antigen I/II family. 10

Method used in this research, therefore, supports and is relevant with the aim of IgY anti *S. mutans* in recognizing both the surface proteins of *S. mutans* with serotype c, d, e, and f, and affinity of its interaction. Thus, the potency of IgY in recognizing Ag I/II specifically shows that IgY used in this research is specific and can be used to restrain Ag I/II of *S. mutans* expressed by all those serotypes. Petersen²⁷ reports that Ag I/II has a good role as adhesin of *S. mutans*, strong tendency to bind saliva component, and a good role as the main initiator in the adhesion and colonization of *S. mutans* in dental pellicle.

The potency of IgY in restraining the surface proteins of *S. mutans* in *in-vitro way*, thus, can be assumed that IgY can prevent the synthesis of glucan from sucrose caused by *S. mutans*, and can decrease the colonization

and carcinogenicity of *S. mutans* in the dental surface. Furthermore, Kruger²⁸ reports that the use of IgY through passive immunization is very effective in protecting caries in rats as the testing animals, which means that IgY can restrain the colonization of *S. mutans* and *S. sobrinus* in *in vivo* experiment. Not only in *S. mutans*, the use of IgY anti *Salmonella enteritis* can also show the good reactivation through ELISA Test,²⁹ Similarly, Rawendra¹¹ reports that IgY can interact well with *Enterophatogenic Escherichia coli* and restrain the colonization of bacteria in the surface of intestine.

Moreover, the research conducted by Smith ⁶ showed that the use of sIgA antibody of *S. mutans* with serotype c, showed that the reactivation of sIgA in the caries free group was higher than that in the caries sensitive group. It means that the antibody can prevent the adhesion of *S. mutans* to hydroxiapatite layers. In other words, the content of sIgA in saliva can prevent the process of caries by restraining the adhesion of *S. mutans*. ³⁰Like sIgA, the result of this *in-vitro* research shows that IgY could be able to recognize Ag I/II considered as the surface proteins of *S. mutans*.

Chismirina³¹ in her research also reports that the use of Polymerase chain reactions (PCR) method successfully determines the serotype of S. mutans strain derived from IPB with serotype d (S. mutans used as the positive control of antigen produced by IgY). Thus, it may be concluded that IgY anti S. mutans with serotype d can recognize proteins of Ag I/II cell surface of S. mutans with serotypes c, d, e, and f similar with S. mutans strain derived from IPB used as the positive control in this research. However, in other to strengthen the result of this research, it needs the further researches about clinical isolate S. mutans strain, especially concerning with the allergic test and cross reaction with some bacteria and fungi of oral cavity, both of which facilitate the adhesion of S. mutans to dental pellicle. Therefore, IgY is expected to be used as biology material in the prevention program of dental caries through passive immunization.

ACKNOWLEDGEMENT

This project was funded by University of Syiah Kuala, Higher Education of Indonesia National Education. National research strategies, contact number 096/H11-P2T/A.01.2009, February, 27 2009 A WELL a BBNAD and Department of Dentistry, Medical Faculty, Syiah Kuala University also molecular Laboratory of FKH IPB and FKG UI.

REFERENCES

- Basri AG, Antonia T, Soeherwin T. Molecular aspect of virulence Streptococcus mutans. Indonesia Journal of Dentistry 2006; 13(2): 107–14
- Samaranayake L. Essential microbiology for dentistry. 3rd ed. Churchill Livingstone Elsevier; 2006. p. 255–93.

- 3. Banas JA. Virulence properties of streptococcus mutans. Frontiers in Bioscience; Journal and Virtual Library 2004; 9: 1267–77.
- Deepan S, Syah H, Roy R, Russell RB. A Novel glucan-binding protein with lipase activity from the oral pathogen Streptococcus mutans. Microbiology 2004; 150: 1947–56.
- Banas JA, Vickerman MM. Glucan binding proteins of the oral streptococci. Crit Rev Oral Biomed2003; 14(2): 89–99.
- Smith DJ. Caries vaccines for the twenty-first century. Journal of Dental Education 2003; 67(10): 1130–7.
- Idone V, Brendtro S, Gillespie R, Kocaj S, Peterson E, Rendi M, Warren W, Michalek S, Krastel K, Cvitkovitch D, Spatafora G. Effect on orphan respon regulator on Streptococcus sucrose-dependent adherence and cariogenesis. Infect Immune 2003; 8(71): 4351–60.
- Biswas S, Biswas I. Role of HtrA in surface protein expression and biofilm formation by Streptococcus mutans. Infection and Immunity 2005; 73 (10): 6923–34.
- Kruger C. Passive immunization against oral phatogens. Division of Clinical Immunology at the Departement of Laboratory Medicine and Center for Oral Biology at NOVUM, Institute of Odontology, Karolinska University Hospital, Huddinge, Stockholm, Sweden. Karolinska University Press; 2004. p. 14–29.
- Crowley PJ, Trevor BS, Isoda R, van Tilburg M, Monika WO, Rebekah AR, William PM, Arnold SB, Brady JL. Requirements for surface expression and function of adhesin P1 from Streptococcus mutans. Infection and Immunity 2008; 76(6): 2456-68.
- Rawendra R. Immunoglobulin Y (IgY) water soluble fraction (WSF) kuning telur kering beku anti Enterophatogenic Escherichia coli (EPEC). Disertation. Sekolah Pascasarjana Institut Pertanian Bogor; 2005. p. 41-100.
- Mitoma M, Oho T, Michibata N, Okano K, Nakano Y, Fukuyama M, Koga T. Passive immunization with bovine milk containing antibody to a cell surface protein antigen-glucosyltransferase fusion protein protects rats against dental caries. Infect Immune 2002; 70(5): 2721–4.
- 13. Chi ZB, Gao YX, Pan Y, Zhang B, Feng XP. The inhibitive effect of IgY toothpaste against oral Streptococcus mutans. Article in Chinese. Department of Preventive & Pediatric Dentistry, Ninth People's Hospital, School of Stomatology, Shanghai Second Medical University. Shanghai 200011. China: Shanghai Kou Qiang Yi Xue; 2004; 13(4): 256–8.
- 14. Wibawan IWT. Pemanfaatan telur ayam sebagai pabrik biologis: produski "yolk immunoglobulin" (IgY) anti plaque dan diare dengan titik berat pada anti Streptococcus mutans, Escherichia coli dan Salmonella enteritidis. Laporan Penelitan. Lembaga Penelitian dan Pemberdayaan Masyarakat IPB, FKH-IPB; 2005. p. 1–20.
- Basri AG, Boy MB, Dewi F. Surface proteins of Streptococcus mutans Ccell that can be recognized by Immunoglobulin Y. Dentika Dental Journal 2006; 11(2): 188–93.
- Yera H, Andiva S, Perret C, Limonne D, Boireau P, Dupovy Canet. Development and evaluation of a Western Blot Kit for diagnosis of human Trichinellosis. American Society for Microbiology; 2003. 10(5): 793–6.

- Carlander D. Avian IgY Antibody In vitro and in vivo. Disertation for the Degree of Doctor of Philosophy (Faculty of Medicine) in Clinical Chemistry presented at Uppsala University; 2002. p. 3–53.
- Zhou Z, Zhou R, Tang Z. Effects of topical application of immunoglobulin yolk on mutans streptococci in dental plaque. Hua Xi Kou Qiang Yi Xue Za Zhi 2003; 21(4): 295–7.
- Wen B, Zhou RJ, Yang J, Tang Z, Zhou Z. The inhibitory effect of anti-streptococcus mutans immunoglobulin of yolk on glucan synthesis of Streptococcus mutans. Shanghai Kou Qiang Yi Xue 2002; 11(2): 141–2.
- Sato Y, Okamoto K, Kagami A, Yamamoto Y, Igarashi T, Kizaki, H. Streptococcus mutans strains harboring collagen-binding adhesion. Research reports. J Dent Res 2004; 87(7): 534–9.
- West AP Jr, Herr AB, Bjorkman PJ. The chicken yolk sac IgY receptor, a functional equivalent of the mammalian MHC-related Fc receptor, is a phospholipase A2 receptor homolog. Immunity 2004; 20(5): 507–8.
- Matsumura M, Izumi T, Matsumoto M, Tsuji M, Fujiwara T, Ooshima T. The role of glucan-binding protein in the cariogenecity of Streptococcus mutans. J Microbiol Immunol 2003; 47(3): 213–15.
- Mattos-Graner RO, Smith DJ. The vaccination approach to control infections leading to dental caries. Bra J Oral Sci 2004; 3(11): 595–608.
- Smith DJ, King WF, Godiska R. Passive transfer of immunoglobulin Y antibody to Streptococcus mutans glucan-binding protein B can confer protection against experimental dental caries. Infect Immun 2001; 69(5): 135–42.
- Mine Y, Kovacs-Nolan J. Chicken egg yolk antibody as therapeutics in enteric infectous disease: a review. J Med Food 2002; 5(3): 159–69.
- Nikki R, Jenny MC, Knneth BT, William PM, Brady JL. William P, McArthur. Characterization of the Streptococcus mutans P1 epitope recognized by immunomodulatory monoclonal antibody 6-11A. Infection and Immunity 2004; 72(8): 4680–88.
- Petersen FC, Assev S, van der Mei HC, Busscher HJ, Scheie AA. Functional variation of the antigen I/II surface protein in Streptococcus mutans and Streptococcus intermedius. Infect Immun 2002; 70: 249–56.
- Kruger C, Pearson SK, Kodama Y, Vacca Smith A, Bowen WH, Hammarstrom L. The effects of egg-derived antibody to glucosyltransferases on dental caries in rats. Caries Res 2004; 8(1): 9–14.
- 29. Lee EN. Anti-bacterial activities of chicken egg yolk antibody (IgY) against enteric pathogens. A thesis Master of Science submitted to the Faculty of Graduate Studies and Research in partial fulfillment of the requirements for the degree of in Food Science and Technology University of Alberta, Canada; 2000. p. 93–106.
- Smith DJ. Dental caries vaccines: prospects and concerns. Crit Rev Oral Biol Med 2002; 13(4): 335–49.
- 31. Chismirina S. Efek Imunoglobulin Y (IgY) sebagai anti adhesin pada pembentukan biofilm oleh Streptococcus mutans (serotype c, e, f) dan Streptococcus sobrinus (serotype d) secara in-vitro. Thesis. Jakarta: Fakultas Kedokeran Gigi Universitas Indonesia; 2006. p. 10–20.